

AD-762 744

MECHANISMS OF REACTIONS OF OXIDIZERS

A. G. Keenan

Miami University

Prepared for:

Office of Naval Research

31 May 1973

DISTRIBUTED BY:

**NTIS**

National Technical Information Service  
U. S. DEPARTMENT OF COMMERCE  
5285 Port Royal Road, Springfield Va. 22151

## **DISCLAIMER NOTICE**

**THIS DOCUMENT IS BEST QUALITY  
PRACTICABLE. THE COPY FURNISHED  
TO DTIC CONTAINED A SIGNIFICANT  
NUMBER OF PAGES WHICH DO NOT  
REPRODUCE LEGIBLY.**

AD 762744

ANNUAL REPORT No. 12

for the period ending: 31 May 1973

Contract Nonr - N00014-67-A-0201-0014


submitted to  
Office of Naval Research  
Material Sciences Division  
Power Program

MECHANISMS OF REACTIONS OF OXIDIZERS

Reproduced by  
NATIONAL TECHNICAL  
INFORMATION SERVICE  
U S Department of Commerce  
Springfield VA 22151

by  
A. G. Keenan

Department of Chemistry  
University of Miami  
Coral Gables, Florida  
33124



Approved for public release  
Distribution Unlimited

None

Security Classification

## DOCUMENT CONTROL DATA - R &amp; D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) University of Miami Department of Chemistry		2a. REPORT SECURITY CLASSIFICATION None	
		2b. GROUP	
3. REPORT TITLE Mechanisms of Reactions of Oxidizers			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Annual Report, June 1, 1972 to May 31, 1973			
5. AUTHOR(S) (First name, middle initial, last name) Keenan, A. G.			
6. REPORT DATE May 31, 1973		7a. TOTAL NO. OF PAGES 8-8	7b. NO. OF REFS 0
8a. CONTRACT OR GRANT NO. N00014-67-A-0201-0014		9a. ORIGINATOR'S REPORT NUMBER(S) Annual Report No. 12	
b. PROJECT NO.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
c.			
d.			
10. DISTRIBUTION STATEMENT Qualified requesters may obtain copies from DDC			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Office of Naval Research Power Program, Code 473 Arlington, VA22217	
13. ABSTRACT Progress during the year in the investigations of the mechanism of decomposition of oxidizers such as ammonium nitrate and ammonium perchlorate is summarized. Areas of investigation include the study of the electrical conductivity of single crystals of ammonium perchlorate, measurement of mobility ratios, ion exchange constants and selectivity ratios of cations in Pyrex immersed in fused nitrates and the study of the electrode reactions at both Cu/CuO and Pt/PtO <sub>2</sub> electrodes in carbonate containing fused nitrates exposed to atmospheres of O <sub>2</sub> , CO <sub>2</sub> , N <sub>2</sub> and H <sub>2</sub> O. Publications issued and pending are listed.			

DD FORM 1473  
1 NOV 65None  
Security Classification

ia

None  
Security Classification

14	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	A ionium Nitrate Perchlorate Propellant Oxidizer EMF Fused Salts Defects Electrical Conductivity						

id

None  
Security Classification

-1-

Annual Report No. 12 - 31 May 1973

Office of Naval Research - Contract N00014-67-A-0201-0014

MECHANISMS OF REACTIONS OF OXIDIZERS

A. G. Keenan

Chemistry Department, University of Miami  
Coral Gables, Florida 33124

Special Report No. 10, comprised of a reprint of the paper "Kinetics of the Low Temperature Thermal Decomposition of Ammonium Perchlorate and Its Catalysis by Copper Ion" by A. G. Keenan and Robert F. Siegmund, Journal of Solid State Chemistry, 4, 362 (1972), was distributed during the year. This work was taken from the Ph.D. Dissertation of R. F. Siegmund.

Special Report No. 11, comprised of a reprint of the paper "The Synergistic Catalysis of Ammonium Nitrate Decomposition - Visible Spectra of Ammine, Chloro and Nitrate Complexes of Copper, Nickel and Cobalt in Fused Ammonium Nitrate", by A. G. Keenan and I. J. Ferrer, Journal of Physical Chemistry, 76, 2844 (1972), was also distributed during the year. This was taken from the M.S. Thesis of Mr. Ferrer.

A publication entitled "Computer Simulation of Defects in Ammonium Perchlorate" by M. Goldstein and A. G. Keenan will appear in the Journal of Solid State Chemistry, 7, No. 3, July (1973). This material is taken from the Ph.D. Dissertation of M. Goldstein. Reprints will be distributed as Special Report No. 12 as soon as available.

A brief manuscript entitled "The Decomposition of Oxalate in Fused Nitrate" by A. G. Keenan and Carlos G. Fernandez has been submitted to the Journal of the Electrochemical Society and a second brief manuscript entitled "The Stability of Carbonate Ion in Fused Nitrate" by A. G. Keenan, Carlos G. Fernandez and Thomas R. Williamson has just been completed.


Research on this project during the year has been in two relatively unrelated areas in the general field of "Mechanisms of Reactions of Oxidizers". One of these deals with the solid state decomposition of ammonium perchlorate, the experimental approach being a study of the electrical conductivity of single crystals both pure and when doped with transition metal ions

or exposed to various atmospheres. The other area concerns an electrometric study of acidity in fused nitrates, with a view to understanding further the mechanism of their decomposition reactions and explaining the synergistic catalytic system discovered earlier.


The work on ammonium perchlorate decomposition follows the earlier theoretical calculations by Dr. Goldstein referenced above. During the year, an enclosed glass system has been built which allows the electrical conductance of single crystals to be measured at temperatures from ambient to 300°C while pumped under high vacuum or exposed to variable pressures of gases such as N<sub>2</sub>, O<sub>2</sub>, HCl, H<sub>2</sub>O and others. Crystals of ammonium perchlorate, pure or doped with various transition metal ions such as Cu and Ni have been grown by crystallization from aqueous solution.

It has been shown that H<sub>2</sub>O and NH<sub>3</sub> individually do not affect the conductivity at room temperature significantly, but when the AP crystal is exposed to both gases in succession a remarkable 15,000% increase in conductivity results. Previous workers have studied the effect of NH<sub>3</sub> but have not apparently noted the significant effect produced when H<sub>2</sub>O and NH<sub>3</sub> are combined. It is believed that some of the oversight may be due to the fact that others have not pumped their AP to the degree of dryness attained in the present work.


The above phenomenon is most readily explained on the basis that the charge carriers in AP are NH<sub>4</sub><sup>+</sup> produced in the experiment described above by the reaction



Combining this with the fact that H<sub>2</sub> is a known cathode product of the conduction process, a reasonable cathode reaction may then be postulated as



A parallel anode reaction which explains the known depression of conductivity by O<sub>2</sub> is



Other work which has also been done includes determining the ratio of surface to bulk conductance by use of a guard ring method. Preliminary results indicate that 99% may be surface. A coulometer has been built to collect and analyze electrode products. A higher voltage than initially estimated will be

required and a power supply for 20,000 volts is now under construction. Conductivity measurements on a pure AP crystal with varying temperature give two very accurate linear regions in  $\log (\sigma T)$  plotted against  $1/T$ . The lower range from 25 to 75° has a correlation coefficient of 0.996 and gives an activation energy of 0.13 eV. The range from 75 to 135° has a correlation coefficient of 0.998 and gives  $E_a$  equal to 1.1 eV.

These experiments will be continued. Ammonium perchlorate doped with various cations and exposed to partial pressures of various known atmospheres, as well as rubidium and cesium perchlorates will be used. The purpose of this research is to elucidate the conduction mechanism and hence discover which ions are mobile and what kinds of defects may be present. Such information in turn should contribute to an understanding of the thermal decomposition mechanism.

In the area of electrometric studies in fused nitrates, research has been conducted along three lines, all of which are ultimately concerned with the study of acidity and its role in the decomposition kinetics of fused oxidizers. First, the principles of operation of Pyrex membrane electrodes have been studied following the method suggested by Conti and Eisenman. This involves measuring voltage/current ratios across the glass membrane while immersed in fused salt. Cation mobility ratios ( $u_i/u_j$ ), ion exchange constants ( $K_{ij}$ ) and selectivity constants ( $k_i$ ) can then be calculated from the data. An alternative procedure which involves electrolyzing the glass to constant conductivity was also developed. The results by the two methods are given in the tables below.

Experimental Values for  $K_{ij}$  and  $u_i/u_j$  in Pyrex Glass  
 $i = \text{Na}^+, j = \text{K}^+ \text{ or } \text{Ag}^+$

T(°C)	325	350			380		413	
	$u_i/u_j$	$K_{ij}$	$u_i/u_j$		$u_i/u_j$		$u_i/u_j$	
$\text{Na}^+ - \text{Ag}^+$	1.22	1.24	0.92	0.86	1.10	0.86	1.0	0.74
$\text{Na}^+ - \text{K}^+$	--	0.257	7.87	--	16.0	50.5	--	82.0
Method	(b)	(a)	(a)	(b)	(a)	(b)	(a)	(b)

(a) by Conti-Eisenman method

(b) electrolysis measurements




Selectivity Ratios for  $\text{Na}^+$ ,  $\text{Ag}^+$  and  $\text{K}^+$  at  $350^\circ\text{C}$  in  
Pyrex Glass

<u>Ion</u>	<u>Selectivity, k</u>
$\text{Na}^+$	1.0
$\text{Ag}^+$	0.78
$\text{K}^+$	0.07


For sodium and silver the results of the two methods agree within expected error. For sodium and potassium the results differ by a factor of nearly four. This is believed to be due to the fact that the composition of the glass changes during the Conti-Eisenman procedure. The mobilities of sodium and potassium are sufficiently different so that the results are influenced by this change, whereas the mobilities of sodium and silver are nearly equal.

The data in the table are not yet complete and further work is required. The data obtained so far in many cases check such scattered literature values as are available. In other cases large discrepancies exist. These will have to be investigated further.

The second line of research in the area of electrometric studies in fused nitrates has dealt with the behavior of a  $\text{Cu}/\text{CuO}/\text{CO}_3^{2-}$  electrode in the presence of  $\text{N}_2$ ,  $\text{O}_2$  and  $\text{CO}_2$  atmospheres, pure and in various mixtures. The results indicate that the electrode behaves reversibly in the presence of  $\text{O}_2$ ,  $\text{CO}_2$  and mixtures of them. The  $E^\circ$  for the electrode at  $350^\circ$  and with  $\text{KNO}_3$  in pure  $\text{O}_2$  is 695 mV while in pure  $\text{CO}_2$  it is 592 mV. When the electrode is subject to mixed gases it behaves according to the respective partial pressures of the gases present. The electrode reaction in accordance with the results so far obtained is:



The mechanism for the electrode reaction may therefore be any of the following three:




This work will be extended to  $\text{NaNO}_3$  and mixtures and is expected to be concluded shortly.

The third line of research in the area of electrometric studies in fused nitrates is similar to the second except that the electrode used is  $\text{Pt/PtO}_2$ . In the course of this work it was necessary to study the stability of  $\text{CO}_3^{2-}$  in fused nitrates because of contradictory statements in the literature. This study was carried out with a pH recorder and constant rate buret to determine the two end points characteristic of carbonate which had been observed previously with colorimetric indicators. Melts were purged with nitrogen or oxygen since these gases have been claimed to remove carbon dioxide from the melt. The results showed no dissociation of carbonate.

It has also been indicated in the literature that addition of dichromate enhances the evolution of carbon dioxide causing the dissociation of carbonate to continue to completion. Results in this laboratory show that while carbon dioxide is evolved, the reaction occurs stoichiometrically as follows:



These results provide clear evidence that the reaction



does not take place in nitrate melts, and thus that carbonate cannot be considered as a source of oxide ion in these melts.

The work on the  $\text{Pt/PtO}_2$  electrode has shown that it behaves in essentially the same manner as the Pt rotating disc electrode previously described by Zamboni. The electrode half-reaction is the same as given earlier for the  $\text{Cu/CuO}$  electrode, when the atmosphere is a mixture of  $\text{CO}_2$  and  $\text{O}_2$ . However, in an atmosphere of  $\text{CO}_2$  only, the reaction is



This has also been proposed by others for a Pt rotating disk electrode. The  $\text{Pt/PtO}_2$  electrode was also found to exhibit non-Nernstian behavior with respect to carbonate in a pure oxygen atmosphere and to be irreversible in a nitrogen atmosphere.